


This article was downloaded by: [98.115.143.186]

On: 09 May 2014, At: 11:48

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of American College Health

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/vach20>

A Randomized Controlled Trial of Koru: A Mindfulness Program for College Students and Other Emerging Adults

Jeffrey M. Greeson PhD^{ab}, Michael K. Juberg BA^b, Margaret Maytan MD^b, Kiera James^c & Holly Rogers MD^{bd}

^a Duke Integrative Medicine, Duke University Medical Center, Durham, North Carolina

^b Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine, Durham, North Carolina

^c Swarthmore College, Swarthmore, Pennsylvania

^d Duke Counseling and Psychological Services, Duke University, Durham, North Carolina

Accepted author version posted online: 05 Feb 2014. Published online: 08 May 2014.

To cite this article: Jeffrey M. Greeson PhD, Michael K. Juberg BA, Margaret Maytan MD, Kiera James & Holly Rogers MD (2014) A Randomized Controlled Trial of Koru: A Mindfulness Program for College Students and Other Emerging Adults, Journal of American College Health, 62:4, 222-233, DOI: [10.1080/07448481.2014.887571](https://doi.org/10.1080/07448481.2014.887571)

To link to this article: <http://dx.doi.org/10.1080/07448481.2014.887571>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Major Article

A Randomized Controlled Trial of Koru: A Mindfulness Program for College Students and Other Emerging Adults

Jeffrey M. Greeson, PhD; Michael K. Juberg, BA; Margaret Maytan, MD; Kiera James; Holly Rogers, MD

Abstract. Objective: To evaluate the effectiveness of Koru, a mindfulness training program for college students and other emerging adults. **Participants:** Ninety students (66% female, 62% white, 71% graduate students) participated between Fall 2012 and Spring 2013. **Methods:** Randomized controlled trial. It was hypothesized that Koru, compared with a wait-list control group, would reduce perceived stress and sleep problems, and increase mindfulness, self-compassion, and gratitude. **Results:** As hypothesized, results showed significant Group (Koru, Wait-List) \times Time (Pre, Post) interactions for improvements in perceived stress ($F[1, 76.40] = 4.50, p = .037, d = .45$), sleep problems ($F[1, 79.49] = 4.71, p = .033, d = .52$), mindfulness ($F[1, 79.09] = 26.80, p < .001, d = .95$), and self-compassion ($F[1, 74.77] = 18.08, p < .001, d = .75$). All significant effects were replicated in the wait-list group. Significant correlations were observed among changes in perceived stress, sleep problems, mindfulness, and self-compassion. **Conclusions:** Results support the effectiveness of the Koru program for emerging adults in the university setting.

Keywords: college health, emerging adulthood, meditation, mindfulness, stress management

According to the 2012 National Survey of College Counseling Centers, 92% of counseling centers are serving increasing numbers of students each year, and 88% of directors report a continued trend towards students presenting with increasing levels of psychopathology.¹ Counseling centers are feeling this strain despite evidence that only a small percentage of struggling students seek treatment from university health centers.^{2,3} The level of stress amongst students is alarming. About 50% of students report significant levels of anxiety and depression,^{2,4} and 16.5% report a history of suicidal or self-harm behavior.² Moreover, sleep disturbance is prevalent in college students and has been proposed as a transdiagnostic risk factor for poor mental health.^{5,6} The effects of mental health problems for students are not insignificant, as students with mental health challenges have difficulties with campus engagement, personal relationships, and graduation rates.⁷ Because of the magnitude of the problem and the limitations on available resources, those working in student health and wellness need creative, developmentally targeted, evidenced-based strategies to effectively reduce stress and improve well-being in student populations. Mindfulness-based interventions may play a role in filling this important need.

Mindfulness, the skill of learning to pay attention, without judgment, to one's present-moment experience, offers a way to improve well-being in diverse populations.⁸ Mindfulness meditation training has been shown to reduce stress and improve mood and academic performance in both college and graduate students.⁹⁻¹² In the United States, mindfulness-based stress reduction (MBSR) and variants of MBSR are the most commonly used methods of teaching mindfulness meditation to adults. MBSR was developed by Jon Kabat-Zinn¹³ and is effective for improving psychological

Dr Greeson and Mr Juberg are with Duke Integrative Medicine at Duke University Medical Center in Durham, North Carolina, and the Department of Psychiatry and Behavioral Sciences at Duke University School of Medicine in Durham, North Carolina. Dr Maytan is with the Department of Psychiatry and Behavioral Sciences at Duke University School of Medicine in Durham, North Carolina. Ms James is with Swarthmore College in Swarthmore, Pennsylvania. Dr Rogers is with the Department of Psychiatry and Behavioral Sciences at Duke University School of Medicine in Durham, North Carolina, and Duke Counseling and Psychological Services at Duke University in Durham, North Carolina.

Copyright © 2014 Taylor & Francis Group, LLC

health, with meta-analyses documenting medium-to-large effect sizes.^{14,15} Although there has been work on developing programs specifically for children and teens,¹⁶ there has been little discussion about targeting mindfulness training specifically towards other critical developmental stages, such as emerging adulthood.

Emerging adulthood is the developmental stage from the late teens through the 20s, which includes most college students.¹⁷ In 2011, 68.2% of graduating high school students enrolled in postsecondary education, and in the fall of 2011 there were nearly 21 million students enrolled in either 2- or 4-year degree-granting institutions.¹⁸ College-attending and non-college-attending emerging adults (EAs) have similar developmental characteristics,¹⁹ and the overall rate of psychiatric disorders is roughly equivalent between college-attending individuals and their non-college-attending peers.⁴ Because the primary emphasis in emerging adulthood is on identity exploration, this life stage is associated with frequent changes and uncertainty in all life spheres.¹⁹

Mindfulness meditation offers a promising tool for managing the stress that may be associated with the challenges of emerging adulthood. However, there are no objective data about what pedagogical methods may be effective for training EAs in mindfulness. Due to their unique developmental stage, EAs may not engage readily in training programs that are designed for older adults due to time constraints, skepticism about the potential benefits, and difficulty maintaining motivation to effect behavior change.²⁰ To address this disparity, Rogers and Maytan²¹ have developed Koru, a manualized training program in mindfulness, meditation, and other mind-body skills, that specifically targets EAs.

Koru is the New Zealand Maori word for the unfurling fern frond, which symbolizes balanced growth. Koru differs from mindfulness programs developed for more general populations of adults in several key ways. First, the program trains EAs in certain mind-body skills, such as abdominal breathing and guided imagery, as well as insight meditation practice. These mind-body skills, for which studies show a positive effect on stress and anxiety,^{22,23} give participants tools for quickly reducing distress, which in turn builds their motivation to continue practicing stress management and self-care in daily life. Second, Koru is relatively brief, with only four 75-minute sessions and 10 minutes of required daily meditation practice. The course is highly structured and attendance at all 4 classes is required. Third, the program is taught in a small-group format to capitalize on peer interactions that EAs value. Fourth, the teaching is very active, addressing skepticism and building motivation quickly. Finally, the language and metaphors used are designed to resonate with the interests and concerns of EAs. In addition to teaching mindfulness skills to cope with stress, Koru also emphasizes cultivating positive emotions such as self-compassion and gratitude. Both compassion and gratitude are considered core qualities of mindfulness,²⁴ and have been associated with lower levels of stress and psychological symptoms and greater well-being in prior studies.^{25,26} Koru has been very popular with students and those who have participated report

positive outcomes on course evaluations.^{20,21} However, until now, Koru has not been subjected to rigorous quantitative evaluation.

The present randomized controlled trial (RCT), therefore, aimed to evaluate the effectiveness of Koru for EAs in a university setting. We hypothesized that college and graduate students who participate in Koru, when compared with a control group, would show reduced stress, improved sleep, and increased mindfulness, self-compassion, and gratitude.

METHODS

Study Design

The RCT had 2 study arms: (1) Koru, the active intervention, and (2) a wait-list control. We chose to use a wait-list control condition to control for the effects of time, because levels of stress, sleep problems, and even self-concept can change during an academic semester. Students who were randomly assigned to the wait-list control group were offered an opportunity to participate in Koru later the same semester. We felt that this design struck a balance between scientific rigor, by benefitting from an experimental design with random assignment to groups, with the use of a culturally normal wait-list process for the Koru program, which should optimize motivation and retention.

Following online informed consent, study participants completed standardized self-report questionnaires via a survey Web site within 1 week before and after participating in the 4-week Koru program. The study protocol was conducted between Fall 2012 and Spring 2013. Full approval by the university's Institutional Review Board was obtained prior to data collection.

Students were eligible for the study if they were (a) currently enrolled as an undergraduate, graduate or professional student; (b) at least 18 years of age; (c) proficient in English; (d) able to use a computer with internet access; and (e) willing to be randomized to either the Koru group or the wait-list control group. Blocked randomization (block size = 2) was used to ensure balanced study groups after every 2 students enrolled. The randomization schedule was created using Research Randomizer (www.randomizer.org) and was performed by the study coordinator who kept group assignments blinded from other study staff who performed statistical analyses. Following randomization, the study coordinator contacted students to inform them of their group assignment. Subsequently, the Koru instructors contacted students enrolled in Koru with an introductory e-mail and a request to begin reading the required text, *Wherever You Go, There You Are* by Jon Kabat-Zinn. To protect against social desirability bias and potential evaluation bias, Koru course instructors were not directly involved in subject recruitment, consent, randomization, or data collection procedures.

Power analyses (G*Power version 3.1.5; Kiel, Germany) based on a difference between 2 independent means (2 groups) at a significance level of .05, a desired power of .80, and a medium-to-large effect size (ie, Cohen's $d = .6$) based on prior meta-analyses, indicated that 45 students per

group would be required to detect a significant difference for the Koru versus wait-list control group.²⁷

Participants

Participants were 90 undergraduate, graduate, and professional students ($M_{\text{age}} = 25.4$, $SD = 5.7$; range = 18–59). Two graduate students, one who was 59 years old and another who was 42, provided baseline survey data but dropped out before attending the Koru group to which they were randomly assigned. The oldest student who provided complete pre- and postsurvey data was 31 years old. Seventy-four students provided data on the post-Koru survey, a response rate of 82%.

As shown in Table 1, the sample comprised mostly women (66%) who were predominantly white (62%) and non-Hispanic (85%). Half of students reported having no religious affiliation or being atheist. Nearly three-quarters were in graduate or professional school. The 2 randomized study groups did not differ on any demographic characteristics, except average hours of sleep per night. The Koru group reported sleeping half an hour less per night, which was statistically significant but still within the normal range of healthy sleep. We defined program completion as attending at least 3 of the 4 Koru classes. There were no significant differences at baseline between students who completed Koru versus those who did not. The only outcome variable that differed significantly was the Isolation subscale of the Self-compassion Scale; the group of students that completed the Koru program reported a mean decrease in their sense of isolation, whereas the group that attended 2 or less Koru classes reported a mean increase in their sense of isolation post intervention.

Procedure

The secure, online survey was administered using Qualtrics software (Provo, Utah) and included basic demographic information and a battery of standardized self-report questionnaires. The questionnaires assessed symptoms of stress, sleep problems, mindfulness of thoughts and feelings, self-compassion, and gratitude. Participants were not offered financial compensation for completing the surveys.

Intervention

As detailed in a recently published treatment manual,²¹ the Koru program is offered at a large university through Counseling and Psychological Services and consists of 4 required 75-minute classes. The classes include 12 to 14 students taught by 1 or 2 teachers. Students typically participate in Koru because they are seeking training in mindfulness and meditation to help reduce stress and enhance psychological well-being. The Koru program is free of charge, voluntary, not graded, and students do not receive course credit for participating. In addition to attending all 4 weekly small-group sessions, the Koru program requires a commitment to practicing meditation for at least 10 minutes daily, completing and returning a daily meditation log, as well as reading the required chapters in the course book, *Wherever You Go, There*

You Are. The meditation log includes space for daily documentation of 2 things for which the student feels grateful. During each class, students learn and practice mindfulness meditation and 1 or 2 mind–body skills, including breathing exercises, walking meditation, guided imagery, and eating meditation.

Measures

Perceived Stress Scale (PSS)

The 10-item Perceived Stress Scale (PSS) measures perceived stress over the past month, defined by the degree to which one feels like life has been unpredictable, uncontrollable, or overwhelming. The PSS scores have demonstrated reliability and construct validity in student samples and EAs.²⁸ Cronbach's α was .83 for PSS scores at both time points.²⁹

Medical Outcome Study Sleep Scale (MOS SLP9)

The sleep scale is one subscale of the Medical Outcomes Study (MOS) health status measure.³⁰ It consists of 12 items to measure 6 sleep dimensions: initiation (time to fall asleep), quantity (hours of sleep each night), maintenance, respiratory problems, perceived adequacy of sleep, and somnolence (daytime sleepiness). The time frame for the responses is "the past 4 weeks." The 9-item summary scale (Sleep Problems Index II) was used for analysis.³⁰ Higher scores indicate greater sleep problems. Cronbach's α was .88 for sleep quality scores at both time points.²⁹

Cognitive and Affective Mindfulness Scale–Revised (CAMS-R)

The 12-item Cognitive and Affective Mindfulness Scale–Revised (CAMS-R) was used to measure 4 aspects of trait mindfulness, including attention, awareness, acceptance, and present-focus. Previous psychometric evaluation of the CAMS-R found scores to be reliable and valid,³¹ and a recent outcome study found that the CAMS-R was sensitive to change following MBSR training.³² Cronbach's α for CAMS-R scores was .88 at baseline and .80 post intervention.²⁹

Self-compassion Scale (SCS)

The 26-item Self-compassion Scale (SCS)³³ was used to assess participants' tendency to treat themselves with kindness and understanding rather than harsh judgment and criticism during difficult times, to perceive difficult experience as part of common humanity, and to consider difficult thoughts and experiences as separate from the core self. The SCS has 6 subscales (Self-judgment, Self-kindness, Isolation, Common Humanity, Mindfulness, Over-identification) that can be summed to attain a total score. SCS scores have been shown to be reliable and valid in previous studies of college students.³³ The SCS was also sensitive to change with a mindful self-compassion (MSC) training program.³⁴ Higher scores on all scales indicate higher levels of self-compassion.

TABLE 1. Baseline Characteristics of Study Participants

Characteristic	Koru group (n = 45)	Wait-list group (n = 45)	Difference statistic
Age			$t(84) = 0.81, p = .423$
<i>M</i> years	25.75	24.76	
<i>SD</i>	6.84	4.15	
Gender			$\chi^2(1) = 0.05, p = .842$
Male	15	16	
Female	30	29	
Ethnicity			See note
Hispanic	4	1	
Non-Hispanic	38	38	
Prefer not to answer	3	5	
Race			$\chi^2(1) = 0.79, p = .375$
White	30	26	
African American	3	2	
Asian American	10	13	
Other	1	3	
Prefer not to answer	1	1	
Religion			$\chi^2(2) = 2.14, p = .343$
No religious affiliation	14	19	
Atheist	5	7	
Buddhist	1	0	
Christian	13	7	
Jewish	1	2	
Hindu	2	1	
Muslim	0	2	
Unitarian/Universalist	1	1	
Other	2	1	
Prefer not to Answer	6	5	
Education			$\chi^2(2) = 0.32, p = .852$
Undergraduate 1st yr	0	3	
Undergraduate 2nd yr	6	2	
Undergraduate 3rd yr	4	3	
Undergraduate 4th yr	2	5	
Undergraduate 5th yr	0	1	
Graduate student	20	20	
Professional Student	13	11	
Sleep			$t(88) = 2.26, p = .027$
<i>M</i> hours per night	7.56	8.09	
<i>SD</i>	1.09	1.16	
Sleep medication			$\chi^2(1) = 0.34, p = .561$
<i>n</i> yes	8	6	
%	18%	13%	

Note. Chi-square could not be computed for ethnicity due to less than 5 Hispanic students per group. Other demographic categories with less than 5 students each were collapsed as follows. Race: white, racial minority. Religion: no religious affiliation, atheist, religious affiliation. Education: undergraduate, graduate, professional student.

Cronbach's α for SCS total scores was .94 at baseline and .95 post intervention.²⁹

Gratitude Questionnaire (GQ-6)

Dispositional gratitude was measured using the GQ-6,³⁵ which has demonstrated good convergent validity with relevant constructs such as positive affect and prosocial traits.³⁶ Cronbach's α for GQ-6 scores was .88 both pre and post intervention.²⁹

Other Measures

Demographic characteristics were assessed at the end of the online survey, and are reported in Table 1.

Data Analysis

All statistics were performed using IBM SPSS version 20 (Armonk, New York). Variables were screened for distributional assumptions prior to analysis. All continuous outcome variables approximated a normal distribution with skewness and kurtosis less than 2.0. Primary analyses of between-group effects were tested using linear mixed-effects models, which provide several statistical advantages over traditional repeated-measures analysis of variance (ANOVA).³⁷ Group assignment was treated as a fixed effect and outcome variables were treated as random effects that could vary within-person and across time. This analytic strategy, based on

restricted maximum likelihood estimation, allowed all subjects to be included in the models, even those with missing data. Therefore, mixed-effects models were consistent with intent-to-treat principles without requiring imputation, such as last observation carried forward, which can bias parameter estimates. Mixed-effects modeling also provided the advantage of accounting for dependence of observations in a repeated-measures design, a phenomenon that frequently violates an assumption of traditional ANOVA using the general linear model.

The primary hypothesis of a significant difference between groups over time was tested using a Group \times Time interaction effect, which indicated the effectiveness of Koru compared with a wait-list control. Within the same model runs, *t* tests were used to test hypotheses about planned contrasts between pre- and postintervention scores within the Koru and control groups, and to test for differences between-groups at baseline and at the postintervention time point. Effect sizes for between-group differences over time were calculated using Cohen's *d* and η^2 .³⁸ Secondary analyses included within-group mixed-effects models to replicate the effects of Koru among wait-listed students who took the program later in the same semester. Secondary analyses also included correlations among changes in outcomes measures as a function of randomized group assignment. Consistent with recommendations for clinical trials, statistical significance for all parameter estimates was set at $z = 1.96$, $\alpha = .05$, 2-tailed.³⁹

RESULTS

Baseline Characteristics and Preliminary Analyses

Baseline levels of perceived stress fell more than 1 standard deviation above the norm for adults between ages 18 and 34 and for adults with an advanced degree.²⁸ Baseline sleep problems fell nearly 1 standard deviation above the general population norm.³⁰ Mindfulness scores fell approximately 1 standard deviation below a published norm for college students,³¹ and was comparable to a sample of adults who participated in an 8-week, self-pay MBSR program.³² Self-compassion fell nearly 1 standard deviation below the college student norm³³ and the mean gratitude score was comparable to the mean reported by scale developers for college students.³⁵ *t* tests revealed no significant differences between groups on primary outcome variables at baseline, except gratitude, which was significantly higher in the Koru group (see Table 2). One subscale of the SCS (Common Humanity) was higher in the Koru group at baseline. Data were relatively normally distributed with all *z* scores for skewness and kurtosis less than 2.0.

Figure 1 provides a flow diagram for recruitment and retention throughout the trial. Overall, 90% of students who enrolled in the study completed the baseline survey and were randomized. Over 82% of randomized students were retained. Of the 45 students initially randomized to Koru, 33% attended all 4 classes, 38% attended 3 or more classes, 18% attended 2 classes, and 4% attended only 1 class. Seven

percent ($n = 3$) did not attend any Koru classes after they were initially assigned to the group. Of the 45 students who participated in Koru after being randomized to the wait-list group, 33% attended all 4 classes, 27% attended 3 or more classes, 11% attended 2 classes, and 2% attended only 1 class. Twenty-seven percent of students ($n = 12$) initially assigned to the wait-list did not attend any Koru classes when they were offered the opportunity later in the semester. Consistent with intent-to-treat, all students who were randomized were analyzed. No adverse events were reported by any participant during the study.

Primary Analyses

As hypothesized, there was a significant Group \times Time interaction for change in perceived stress scores (see Table 2). Whereas perceived stress scores for the control group did not change over time ($t = 0.71$, $p = .48$), perceived stress decreased significantly in the Koru group ($t = 3.62$, $p = .001$). A medium effect size on perceived stress was observed for Koru compared with the wait-list control (Table 2), as evidenced by a half standard deviation larger reduction in perceived stress scores ($d = .45$), with 5% of the variance in perceived stress scores over time explained as a function of group assignment ($\eta^2 = .048$).

There was also a significant Group \times Time interaction for change in sleep problems (see Table 2). Whereas sleep quality scores for the control group did not change over time ($t = 0.03$, $p = .98$), overall sleep quality improved significantly in the Koru group ($t = 3.04$, $p = .003$). A medium effect size on sleep problems was observed for Koru compared with the wait list control (Table 2), as evidenced by a half-standard deviation larger reduction in sleep quality scores ($d = .52$), with 6% of the variance in sleep quality scores over time explained as a function of group assignment ($\eta^2 = .063$).


A significant Group \times Time interaction was observed for change in mindfulness scores (see Table 2). Whereas mindfulness scores for the control group did not change over time ($t = 0.59$, $p = .56$), mindfulness increased significantly in the Koru group ($t = -6.60$, $p < .001$). A large effect size on mindfulness of thoughts and feelings was observed for Koru compared with the wait-list control (Table 2), as evidenced by a 1 standard deviation larger increase in CAMS-R mindfulness scores ($d = -.95$), with 18% of the variance in mindfulness scores over time explained as a function of group assignment ($\eta^2 = .184$).

A significant Group \times Time interaction was also observed for change in self-compassion scores (see Table 2). Whereas self-compassion total scores for the control group did not change over time ($t = -0.20$, $p = .84$), they improved significantly in the Koru group ($t = -6.38$, $p < .001$). A large effect size on total self-compassion scores was observed for Koru compared with the wait-list control (Table 2), as evidenced by a three-quarter standard deviation larger increase in total self-compassion scores ($d = -.75$), with 12% of the variance in total self-compassion scores over time explained as a function of group assignment ($\eta^2 = .123$).

TABLE 2. Changes in Outcome Variables for Koru and Wait-List Control Groups

Outcome	Koru group				Wait-list control				Mixed-model ANOVA (Group × Time)			
	Pre		Post		Pre		Post		F	p	d	η^2
	M	SD	M	SD	M	SD	M	SD				
Perceived stress	23.71	5.65	20.43*	6.20	25.66	5.65	25.04	5.96	76.40	.04	.45	.048
Sleep problems	41.36	16.21	32.52*	18.06	42.91	16.21	42.84	17.23	79.49	.03	.52	.063
Mindfulness	28.69	5.63	33.77*	6.06	27.22	5.63	26.79	5.87	79.09	.00	-.95	.184
Self-compassion (total)	2.73	0.66	3.25*	0.70	2.61	0.66	2.62	0.68	20.07	.00	-.75	.123
Self-kindness	2.64	0.77	3.20*	0.84	2.47	0.77	2.55	0.80	9.20	.00	-.06	.083
Common Humanity	2.79	0.84	3.14*	0.90	2.41	0.84	2.65†	0.87	0.51	.48	-.13	.004
Mindfulness	3.07	0.76	3.46*	0.82	2.86	0.76	2.85	0.80	8.03	.01	-.51	.061
Self-judgment	2.67	0.79	3.28*	0.87	2.74	0.79	2.64	0.83	15.39	.00	-.86	.156
Isolation	2.54	0.88	3.16*	0.95	2.55	0.88	2.46	0.91	17.56	.00	-.79	.135
Over-identification	2.68	0.85	3.30*	0.91	2.61	0.85	2.60	0.88	16.81	.00	-.72	.115
Gratitude	35.93	6.96	36.32	7.51	32.58	6.96	32.31	7.26	0.24	.62	-.09	.002

Note. Pre and post scores from mixed-effects models are shown as estimated marginal means (*SD*). Statistics are shown for Group × Time effect, which tested for the effectiveness of Koru over time compared with the wait-list control. Cohen's *d* = difference between pre-post change in Koru and pre-post change in wait-list divided by pooled *SD*. Range: .20 = small, .50 = medium, .80 = large.³⁸ η^2 range for percent variance explained in outcome due to group assignment: .01 = small, .06 = medium, .14 = large.³⁸ Higher scores for all self-compassion scales indicate higher levels of self-compassion. **p* < .005; †*p* < .05 for within-group changes by paired *t* test (see text).



To examine the sources of change in overall levels of self-compassion, we ran separate mixed-effects models on each of the self-compassion subscales (see Table 2). Whereas the wait-list control group only reported a significant pre–post change on 1 subscale, Common Humanity, the Koru group reported significant pre–post changes on all 6 subscales: increased Self-kindness, Common Humanity, and Mindfulness, and decreased Self-judgment, Isolation, and Over-identification (see Table 2). Effect sizes

for SCS subscales fell in the medium range for Self-kindness and Mindfulness, and in the large range for Self-judgment, Isolation, and Over-identification. The Common Humanity subscale, which improved significantly for both groups, showed a small effect size in favor of Koru (see Table 2).

The Group × Time interaction for change in gratitude was not statistically significant. Neither group reported a significant pre–post change in gratitude (see Table 2).

Secondary Analyses

Mixed-effects models on students originally assigned to the wait list ($n = 45$) who took Koru later in the semester replicated all significant effects observed among students who were randomized to take Koru first. Within-subjects analyses modeled change over 3 time points: baseline, post wait-list, and post Koru. Significant effects of time (all p values $< .005$) were accounted for by significant changes from time 2 to time 3 for the following variables: PSS ($t = 3.56$, $p = .001$), sleep problems ($t = 12.75$, $p < .001$), mindfulness ($t = 5.08$, $p < .001$), self-compassion total ($t = 6.24$, $p < .001$), self-judgment ($t = 4.37$, $p < .001$), isolation ($t = 4.70$, $p < .001$), over-identification ($t = 4.10$, $p < .001$), self-kindness ($t = 3.68$, $p < .001$), and mindfulness ($t = 4.14$, $p < .001$). Consistent with students randomized to take Koru first, students randomized to take Koru later in the semester had large effects sizes (within-subjects Cohen's $d = t/\sqrt{n}$) for changes in mindfulness ($d = .76$) and total self-compassion scores ($d = .93$), and medium effect sizes for perceived stress ($d = .53$) and most self-compassion subscales ($d = .55-.70$). The effect size for improvement in sleep problems was substantially larger ($d = 1.90$), and the effect on an increased sense of common humanity was marginally significant ($t = 1.96$, $p = .054$) with a small effect size ($d = .29$).

As expected, numerous significant correlations were observed between changes in perceived stress, sleep problems, mindfulness, and self-compassion among students originally randomized to Koru (see Table 3). Notably, changes in mindfulness, self-compassion total scores, self-judgment, and over-identification were significantly correlated with all other outcome variables (except gratitude) in the Koru group. In contrast, compared with students randomly assigned to Koru, students assigned to the wait-list group reported fewer than half as many significant correlations (see Table 3). These findings are consistent with the hypothesis that improvements in stress, sleep, mindfulness, and compassion are interrelated during Koru.

COMMENT

Emerging adulthood, the transitional stage between high school and adulthood, has been defined as a distinct developmental stage with unique needs and challenges.^{17,19} This age group struggles with concerning levels of stress-related psychological conditions, including anxiety,⁴⁰ depression,² and sleep disturbance.⁵ Mindfulness-based interventions have proved effective for reducing stress in university students.⁴¹ The main results from this RCT found that Koru, a new mindfulness training program designed specifically for EAs, produced significant improvements in perceived stress, sleep quality, mindfulness, and self-compassion in a university student cohort of EAs. Gratitude was not measurably affected. Koru was well liked by students as reflected by high enrollment and retention in the study.

The results of this trial replicate and extend previous studies of meditation and mindfulness training for university students. For example, the significant effect on lowering per-

ceived stress was similar in magnitude to a published RCT on college students that compared the effect of 2 meditation programs with a wait-list control group.⁹ Lower perceived stress following participation in Koru is also consistent with another wait-list controlled trial with medical students.¹⁰ Other RCTs that found significant reductions in perceived stress and psychological distress included a 6-week mind-body program based on the relaxation response and cognitive-behavioral therapy skills⁴² and a 3-arm trial that compared 5 sessions of mindfulness meditation training, somatic relaxation training, and a wait-list control.⁴³ Thus, the significant stress reducing effect of Koru in this study was comparable to other, often longer, stress management and skill building programs for students.

Koru also produced a significant decrease in sleep problems, an effect that has previously been limited to uncontrolled studies.⁴⁴ Whereas RCTs of MBSR and mindfulness-based cognitive therapy (MBCT) have recently documented improvements in objective measures of sleep quality among adult populations with insomnia⁴⁵ and depression,⁴⁶ the present study adds to the literature by demonstrating that improvement in subjective sleep quality among university students is directly correlated with increased levels of mindfulness and self-compassion.

Consistent with a recent meta-analysis of over 12,000 participants in mindfulness-based therapies, Koru produced clinically significant effect sizes on multiple measures of psychological well-being.¹⁴ The largest effect sizes in this study were seen for increases in mindfulness and self-compassion. The increase in mindfulness with Koru was comparable to increases in mindfulness seen with an 8-week MBSR program,³² an 8-week MBCT program,⁴⁷ and an 8-week MSC program.³⁴ Although Koru's relatively large effect size for self-compassion ($d = .75$) was substantially less than that for MSC ($d = 1.67$), it had somewhat stronger effects on mindfulness and stress. This difference may reflect that Koru does not specifically target self-compassion, but rather mindful attention, awareness, relaxation, and self-regulation. Nevertheless, changes in self-compassion in this study were comparable to those seen in other primary mindfulness interventions such as MBSR.⁴⁸ Given the association between higher levels of self-compassion and better psychological well-being,^{25,34} this study suggests that Koru may promote student health, in part, by helping EAs develop a healthier way of relating to themselves in difficult times—a way marked by greater self-kindness and less self-judgment; a greater sense of connection through common humanity and less isolation; and better emotional balance through greater mindfulness and less over-identification with one's problems.

Secondary findings demonstrated that all of the significant effects observed in a group of students randomized to receive Koru could be replicated in wait-listed students later in the same semester. Moreover, correlations among improvements in perceived stress, sleep problems, mindfulness, and self-compassion were predominantly significant in the Koru group. This finding indicates that Koru-related benefits in mental and behavioral health are interdependent.

TABLE 3. Correlations Among Study Variables for Koru Group (n = 35) Below Diagonal and Wait-List Control (n = 39) Above Diagonal

	1	2	3	4	5	6	7	8	9	10	11
1. Perceived stress (PSS)	—	.322*	-.142	.199	-.094	.161	.241	-.275†	-.267	-.243	-.164
2. Sleep problems (MOS)	.313†	—	.059	-.250	-.238	-.021	-.019	-.138	.013	-.334*	.100
3. Mindfulness (CAMS)	-.611***	-.450**	—	.496**	.454**	.123	.361*	.187	.098	.315†	.057
4. Self-compassion (total)	-.459**	-.405*	.711**	—	.661**	.060	.565**	.731**	.426**	.536**	.239
5. Self-kindness (SCS)	-.322†	-.335*	.613**	.821**	—	.039	.537**	.253	-.096	.207	-.070
6. Common Humanity (SCS)	-.243	-.309†	.523**	.628**	.416*	—	.191	-.354*	-.145	-.192	-.277†
7. Mindfulness (SCS)	-.255	-.262	.594**	.799**	.657**	.553**	—	.145	-.052	.024	-.025
8. Self-judgment (SCS)	-.486**	-.334*	.582**	.881**	.642**	.487**	.568**	—	.456**	.459**	.417**
9. Isolation (SCS)	-.388*	-.196	.462**	.642**	.396*	.077	.400*	.563**	—	.048	.266
10. Over-identification (SCS)	-.378*	-.444**	.505**	.825**	.624*	.427*	.593**	.691**	.508**	—	.293†
11. Gratitude (GQ-6)	-.203	-.286†	.276	.241	.143	.329†	.118	.222	.174	.117	—

Note. PSS = Perceived Stress Scale; MOS = Medical Outcomes Study; CAMS = Cognitive and Affective Mindfulness Scale; SCS = Self-compassion Scale; GQ-6 = Gratitude Questionnaire.
 †p < .10; *p < .05; **p < .01; ***p < .001.

Taken together, the significant changes in perceived stress, sleep, mindfulness and self-compassion with Koru suggest that tailoring mindfulness training to emerging adulthood by incorporating other mind–body skills, using fewer class sessions, requiring shorter but mandatory home practice as well as a course text, and contextualizing mindfulness training to reflect common interests and concerns of EAs is an effective strategy.

The relative brevity of the training model may contribute to the high retention rates and thus effectiveness of Koru. Students identify time pressures as one of the primary reasons for not seeking help for mental health concerns.³ Thus, Koru was designed to be as efficient as possible. Koru involves 5 hours of class time over 4 weeks, plus the expectation of 10 minutes of practice/day. In comparison, MBSR involves 27 hours of class time over 8 weeks, plus the expectation of 45 minutes/day of meditation practice.⁴⁹ The decreased time requirement of Koru appears to work well for young adults who are juggling competing pressures. The present study demonstrates that significant results can be achieved within a relatively short, 4-week time frame. Others have found benefits with brief exposure to mindfulness training. For example, one 5-session training model that included a daylong meditation retreat found significant decreases in psychological distress and rumination, and significant increases in positive mood, among undergraduate and graduate students reporting distress.⁴³ Another recent study found reductions in anxiety and associated neural correlates with only 4 20-minute mindfulness trainings.⁵⁰ The relatively high retention rate of 82% in our study may reflect the appeal of Koru for EAs.

Strengths and Limitations

The strengths of the study included a randomized experimental design, adequate statistical power provided by a relatively large sample size, inclusion of all available data from all randomized subjects, and relatively high levels of program adherence and survey completion. Limitations included reliance on self-report surveys for primary outcome measures, which are susceptible to recall bias; limited diversity in the student sample, which precludes broader generalizability; and use of pre- and postintervention measures, which limits understanding of when beneficial effects started to occur and how long beneficial effects might last after Koru training. Future research can build on the present study by using an objective measure of mindfulness, such as monitoring breaths or heart beat while practicing meditation,⁵¹ enrolling a more diverse group of students, and including both interim and follow-up assessments to better describe the process and durability of change following Koru.

Conclusion

College students make up a substantial fraction of the EA demographic.¹⁸ A large proportion of college students report significant stress-related mental health concerns, including anxiety and depression.⁴¹ These types of mental health concerns impact students both interpersonally and

academically.⁷ Mindfulness-based interventions are known to provide significant improvement in anxiety, depression, and stress reactivity in adults and university student populations.^{41,52} Only a small percentage of struggling students receive treatment from university health centers^{2,3}; therefore, it is critical that a broader range of mental-health outreach services be provided for students. The results of this study show that Koru, a relatively brief, developmentally targeted mindfulness program, was effective at reducing symptoms of stress, enhancing psychological well-being, and promoting sleep—a key health behavior—for EAs who are students in a university setting. These findings support Koru as a viable intervention for student counseling centers and other agencies that seek to provide cost-effective, low-stigma interventions for students suffering from unmanageable levels of stress. Further research is needed to explore the effectiveness of Koru in other EA populations in community colleges, historically black colleges and universities, and nonacademic settings.

ACKNOWLEDGMENTS

We thank the staff at Duke Counseling and Psychological Services, including the director of outreach services Gary Glass, PhD, and the administrative assistant for outreach services, Mary Cicinnati. We also thank all of the undergraduate, graduate, and professional students who volunteered their valuable time to participate in the trial and complete our survey. Finally, we thank Karen Caldwell, PhD, for assistance with SPSS code for the mixed-effects models, and Dennis Carmody, BA, for technical assistance with randomization procedures and database management.

FUNDING

This work was supported by grant R00 AT004945 from the National Center for Complementary and Alternative Medicine (NCCAM) to JMG. The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health (NIH).

CONFLICT OF INTEREST DISCLOSURE

After completing the current research study, Dr Rogers co-founded the Koru Center for Mindfulness, of which she has an ownership share; if Koru is found to be an effective intervention, she could benefit financially. Dr Maytan has received compensation for teaching from the Koru Center for Mindfulness. Dr Greeson, Mr Juberg, and Ms James, who were responsible for collecting and analyzing the data for this study, report no financial relationship relevant to the subject of this article. The authors confirm that the research presented in this article met the ethical guidelines, including adherence to the legal requirements, of the United States and received approval from the Institutional Review Board of Duke University.

NOTE

For comments and further information, address correspondence to Jeffrey M. Greeson, PhD, Duke Integrative Medicine, Duke University Medical Center, DUMC Box 102904, Durham, NC 27710, USA (e-mail: jeffrey.greeson@duke.edu; jeffreymgreeson@gmail.com).

REFERENCES

- Gallagher RP. *National Survey of Counseling Center Directors*. Alexandria, VA: International Association of Counseling Services; 2004.
- Garlow SJ, Rosenberg J, Moore JD, et al. Depression, desperation, and suicidal ideation in college students: results from the American foundation for suicide prevention college screening project at Emory University. *Depress Anxiety*. 2008;25:482–488.
- Downs MF, Eisenberg D. Help seeking and treatment use among suicidal college students. *J Am Coll Health*. 2012;60:104–114.
- Blanco C, Okuda M, Wright C, et al. Mental health of college students and their non-college-attending peers: results from the national epidemiologic study on alcohol and related conditions. *Arch Gen Psychiatry*. 2008;65:1429–1437.
- Gaultney JF. The prevalence of sleep disorders in college students: impact on academic performance. *J Am Coll Health*. 2010;59:91–97.
- Harvey AG. Insomnia, psychiatric disorders, and the transdiagnostic perspective. *Curr Dir Psychol Sci*. 2008;17:299–303.
- Salzer MS. A comparative study of campus experiences of college students with mental illnesses versus a general college sample. *J Am Coll Health*. 2012;60:1–7.
- Keng S, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: a review of empirical studies. *Clin Psychol Rev*. 2011;31:1041–1056.
- Oman D, Shapiro SL, Thoresen CE, Plante TG, Flinders T. Meditation lowers stress and supports forgiveness among college students: a randomized controlled trial. *J Am Coll Health*. 2008;56:569–578.
- Warnecke E, Quinn S, Ogden K, Towle N, Nelson MR. A randomised controlled trial of the effects of mindfulness practice on medical student stress levels. *Med Educ*. 2011;45:381–388.
- Franco C, Mañas I, Cangas AJ, et al. The applications of mindfulness with students of secondary school: results on the academic performance, self-concept and anxiety. In: Lytras MD, Ordonez De Pablos P, Ziderman A, et al, eds. *Knowledge Management, Information Systems, E-Learning, and Sustainability Research*. Berlin, Germany: Springer-Verlag; 2010:83–97.
- Mrazek MD, Franklin MS, Phillips DT, Baird B, Schooler JW. Mindfulness training improves working memory capacity and GRE performance while reducing mind wandering. *Psychol Sci*. 2013;24:776–781.
- Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: theoretical considerations and preliminary results. *Gen Hosp Psychiatry*. 1982;4:33–47.
- Khoury B, Lecomte T, Fortin G, et al. Mindfulness-based therapy: a comprehensive meta-analysis. *Clin Psychol Rev*. 2013;33:763–771.
- Eberth J, Sedlmeier P. The effects of mindfulness meditation: a meta-analysis. *Mindfulness*. 2012;3:174–189.
- Burke CA. Mindfulness-based approaches with children and adolescents: a preliminary review of current research in an emergent field. *J Child Fam Stud*. 2010;19:133–144.
- Arnett JJ. Emerging adulthood: a theory of development from the late teens through the twenties. *Am Psychol*. 2000;55:469–480.
- Aud S, Hussar W, Johnson F, et al. *The Condition of Education 2012*. Washington, DC: National Center for Education Statistics, US Department of Education; 2012.
- Arnett JJ. *Emerging Adulthood: The Winding Road From the Late Teens Through the Twenties*. New York, NY: Oxford University Press; 2004.
- Rogers HB. Koru: teaching mindfulness to emerging adults. *New Direct Teach Learn*. 2013;2013(134):73–81.
- Rogers H, Maytan M. *Mindfulness for the Next Generation: Helping Emerging Adults Manage Stress and Lead Healthier Lives*. New York, NY: Oxford University Press; 2012.
- Finkelstein C, Brownstein A, Scott C, Lan Y. Anxiety and stress reduction in medical education: an intervention. *Med Educ*. 2007;41:258–264.
- Saunders PA, Tractenberg RE, Chaterji R, et al. Promoting self-awareness and reflection through an experiential mind-body skills course for first year medical students. *Med Teach*. 2007;29:778–784.
- Shapiro S, Schwartz G, Santerre C. Meditation and positive psychology. In: Snyder CR, Lopez SJ, eds. *Handbook of Positive Psychology*. New York, NY: Oxford University Press; 2002:632–645.
- Neff KD, Rude SS, Kirkpatrick KL. An examination of self-compassion in relation to positive psychological functioning and personality traits. *J Res Pers*. 2007;41:908–916.
- Fredrickson BL, Cohn MA, Coffey KA, Pek J, Finkel SM. Open hearts build lives: positive emotions, induced through loving-kindness meditation, build consequential personal resources. *J Pers Soc Psychol*. 2008;95:1045–1062. doi: 10.1037/a0013262.
- Faul F, Erdfelder E, Lang A, Buchner A. G* power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39:175–191.
- Cohen S, Janicki-Deverts D. Who's stressed? Distributions of psychological stress in the united states in probability samples from 1983, 2006, and 2009. *J Appl Soc Psychol*. 2012;42:1320–1334.
- Cronbach, L. J. Coefficient alpha and the internal structure of tests. *Psychometrika*. 1951;16:297–334.
- Spritzer KL, Hays RD. *MOS Sleep Scale: A Manual for Use and Scoring. Version 1.0*. Los Angeles, CA: RAND; November 2003.
- Feldman G, Hayes A, Kumar S, Greeson J, Laurenceau JP. Mindfulness and emotion regulation: the development and initial validation of the cognitive and affective mindfulness scale-revised (CAMS-R). *J Psychopathol Behav Assess*. 2007;29:177–190.
- Greeson JM, Webber DM, Smoski MJ, et al. Changes in spirituality partly explain health-related quality of life outcomes after mindfulness-based stress reduction. *J Behav Med*. 2011;34:508–518.
- Neff KD. The development and validation of a scale to measure self-compassion. *Self Identity*. 2003;2:223–250.
- Neff KD, Germer CK. A pilot study and randomized controlled trial of the Mindful Self-compassion program. *J Clin Psychol*. 2013;69:28–44.
- McCullough ME, Emmons RA, Tsang JA. The grateful disposition: a conceptual and empirical topography. *J Pers Soc Psychol*. 2002;82:112–27.
- McCullough ME, Tsang J, Emmons RA. Gratitude in intermediate affective terrain: links of grateful moods to individual differences and daily emotional experience. *J Pers Soc Psychol*. 2004;86:295–309.
- Singer JD, Willett JB. *Applied Longitudinal Data Analysis: Modeling Change and Event Occurrence*. New York, NY: Oxford University Press; 2003.
- Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum; 1988.

39. Friedman LM, Furberg CD, DeMets DL. *Fundamentals of Clinical Trials*. New York, NY: Springer; 1998.
40. Keyes CL, Eisenberg D, Perry GS, Dube SR, Kroenke K, Dhingra SS. The relationship of level of positive mental health with current mental disorders in predicting suicidal behavior and academic impairment in college students. *J Am Coll Health*. 2012;60:126–133.
41. Regehr C, Glancy D, Pitts A. Interventions to reduce stress in university students: a review and meta-analysis. *J Affect Disord*. 2013;148:1–11.
42. Deckro GR, Ballinger KM, Hoyt M, et al. The evaluation of a mind/body intervention to reduce psychological distress and perceived stress in college students. *J Am Coll Health*. 2002;50:281–287.
43. Jain S, Shapiro SL, Swanick S, et al. A randomized controlled trial of mindfulness meditation versus relaxation training: effects on distress, positive states of mind, rumination, and distraction. *Ann Behav Med*. 2007;33:11–21.
44. Winbush NY, Gross CR, Kreitzer MJ. The effects of mindfulness-based stress reduction on sleep disturbance: a systematic review. *Explore (NY)*. 2007;3:585–591.
45. Gross CR, Kreitzer MJ, Reilly-Spong M, et al. Mindfulness-based stress reduction versus pharmacotherapy for chronic primary insomnia: a randomized controlled clinical trial. *Explore (NY)*. 2011;7:76–87.
46. Britton WB, Haynes PL, Fridel KW, Bootzin RR. Polysomnographic and subjective profiles of sleep continuity before and after mindfulness-based cognitive therapy in partially remitted depression. *Psychosom Med*. 2010;72:539–548.
47. Kuyken W, Watkins E, Holden E, et al. How does mindfulness-based cognitive therapy work? *Behav Res Ther*. 2010;48:1105–1112.
48. Birnie K, Speca M, Carlson LE. Exploring self-compassion and empathy in the context of mindfulness-based stress reduction (MBSR). *Stress Health*. 2010;26:359–371.
49. Kabat-Zinn J. *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. New York, NY: Delacorte Press; 1990.
50. Zeidan F, Martucci KT, Kraft RA, McHaffie JG, Coghill RC. Neural correlates of mindfulness meditation-related anxiety relief [published online ahead of print April 24, 2013]. *Soc Cogn Affect Neurosci*. doi:10.1093/scan/nst041.
51. Sauer S, Walach H, Schmidt S, et al. Assessment of mindfulness: review on state of the art. *Mindfulness*. 2013;4:3–17.
52. Hofmann SG, Sawyer AT, Witt AA, Oh D. The effects of mindfulness-based therapy on anxiety and depression: a meta-analytic review. *J Consult Clin Psychol*. 2010;78:169–183.

Received: 22 June 2013
 Revised: 12 January 2014
 Accepted: 20 January 2014