

There is a great deal of evidence to prove that immunization of children does more harm than good. - Dr. J. Anthony Morris (formerly Chief Vaccine Control Officer at the US Federal Drug Admin.)

"My suspicion, which is shared by others in my profession, is that the nearly 10,000 SIDS deaths that occur in the United States each year are related to one or more of the vaccines that are routinely given children. The pertussis vaccine is the most likely villain, but it could also be one or more of the others." --Dr. Mendelsohn, M.D.

"The evidence for indicting immunizations for SIDS is circumstantial, but compelling. However, the keepers of the keys to medical-research funds are not interested in researching this very important lead to the cause of an ongoing, and possibly preventable, tragedy. Anything that implies that immunizations are not the greatest medical advance in the history of public health is ignored or ridiculed. Can you imagine the economic and political import of discovering that immunizations are killing thousands of babies?" --Dr. Douglass M.D.

"Crib death was so infrequent in the pre-vaccination era that it was not even mentioned in the statistics, but it started to climb in the 1950s with the spread of mass vaccination against diseases of childhood. -- Harris L.Coulter, PhD."

"There is **insufficient** evidence to support routine vaccination of healthy persons of any age. - Paul Frame, M.D., Journal of Family Practice"

"Delay of DPT immunization until 2 years of age in Japan has resulted in a dramatic decline in adverse side effects. In the period of 1970-1974, when DPT vaccination was begun at 3 to 5 months of age, the Japanese national compensation system paid out claims for 57 permanent severe damage vaccine cases, and 37 deaths. During the ensuing six year period 1975-1980, when DPT injections were delayed to 24 months of age, severe reactions from the vaccine were reduced to a total of eight with three deaths. This represents an 85 to 90 percent reduction in severe cases of damage and death." --Raymond Obomsawin, M.D

"Aluminum is an experimentally **demonstrated neurotoxin** and **the most commonly used vaccine adjuvant**. Despite almost 90 years of widespread use of aluminum adjuvants, medical science's understanding about their mechanisms of action **is still remarkably poor**. There is also a concerning **scarcity of data** on toxicology and pharmacokinetics of these compounds. In spite of this, the notion that aluminum in vaccines is safe appears to be widely accepted. Experimental research, however, **clearly shows that aluminum adjuvants have a potential to induce serious immunological disorders in humans**. In particular, aluminum in adjuvant form **carries a risk for autoimmunity, long-term brain inflammation and associated neurological complications** and may thus have profound and widespread adverse health consequences. In our opinion, the **possibility that vaccine benefits may have been overrated and the risk of potential adverse effects underestimated**, has not been rigorously evaluated in the medical and scientific community"
<http://www.ncbi.nlm.nih.gov/pubmed/21568886>

BE AWARE OF THE DELAYED REACTION!

HOW LONG AFTER A VACCINE DO INJURIES OCCUR?

**IMMEDIATELY TO YEARS LATER.
OFTEN 13-19 DAYS AFTER VACCINATION.**

YOU KNOW THE SCREAMS

Arched back. Inconsolable crying. ENCEPHALITIS. BRAIN DAMAGE. Many parents do not recognize the link between vaccination and reaction or injury because often times the reaction is delayed until after the vaccine virus incubates. Almost a month after they get their shots. **This takes up to and around 3-4 weeks after the child is injected to occur.** WATCH YOUR BABY AROUND THE 2ND-3RD WEEK! KEEP THEM CLOSE TO YOU! COMFORT THEM IF THEY BECOME SICK. NO TYLENOL OR ADVIL. THIS HURTS THE DETOXIFICATION PROCESS AND PUTS YOUR BABY AT GREATER RISK FOR SEIZURES.

"Human tumors containing SV-40 cancer virus are now found in 61% of autopsies of those who died after age 55. These are linked directly to the SV-40 virus contained in the polio vaccines given from the 1950's through the 1960's."

<http://ppig.me/2011/05/13/vaccineshuman-and-animal-dna/>




EMAIL/FACEBOOK
Educate4theinjured@hot-shot.com
 For FREE BROCHURES

www.Educate4theInjured.org


EDUCATE

BEFORE YOU VACCINATE!

"THUS VACCINATION DOES NOT ACCOUNT FOR THE IMPRESSIVE DECLINES IN MORTALITY SEEN IN THE FIRST HALF OF THE CENTURY"




Scarlet Fever



Annual Summary of Vital Statistics: Trends in the Health of Americans During the 20th Century
 Guyer, et al.
 American Academy of Pediatrics 2000; 106:6 1307-1317

Measles



SOURCE: Data - Official Year Books of the Commonwealth of Australia, as reproduced in Greg Beattie's book "Vaccination A Parent's Dilemma"

Whats **INSIDE** the



by the age of 6?

26 by the 1st yr of life

Research the connections.

"A single vaccine given to a six-pound newborn is the equivalent of giving a 180-pound adult 30 vaccinations on the same day."

Dr. Boyd Haley, Professor and Chair, Dept. of Chemistry, University of Kentucky (2001)



How does the child's body process these ingredients?

Ammonium Sulfate: EDF Suspected - gastrointestinal or liver toxicant
neurotoxicant
respiratory toxicant

Amphotericin B: MME definition - "a drug used to treat fungus infections. Known allergy to this drug prohibits use. Side effects include blood clots, blood defects, kidney problems, nausea and fever. When used on the skin, allergic reactions can occur."

Aluminum: EDF Suspected - cardiovascular or blood toxicant
neurotoxicant
respiratory toxicant
More hazardous than most chemicals in 2 out of 6 ranking systems
On at least 2 federal regulatory lists

Beta-Propiolactone: EDF Recognized - carcinogen
EDF Suspected - gastrointestinal or liver toxicant
respiratory toxicant
skin or sense organ toxicant
More hazardous than most chemicals in 3 out of 3 ranking systems
On at least 5 federal regulatory lists
Ranked as one of the most hazardous compounds (worst 10%) to humans

Thimerosal: ethyl mercury
EDF Recognized - development toxicant
Suspected - skin or sense organ toxicant

Sucrose: refined sugar

Gentamicin Sulfate: an antibiotic

Hydrolyzed Gelatin: obtained from selected pieces of calf and cattle skins, de-mineralized cattle bones (ossein) and porkskin

Neomycin: an antibiotic

Polyribosylribitol: a component of the Hib bacterium

Polysorbate: EDF Suspected - skin or sense organ toxicant

Polymyxin: an antibiotic

Tri(n)butylphosphate:
EDF Suspected - kidney toxicant
neurotoxicant
More hazardous than most chemicals in 2 out of 3 ranking systems
On at least 1 federal regulatory list

Streptomycin: an antibiotic

Phenol: aka Carbohic Acid
EDF Suspected - cardiovascular or blood toxicant
developmental toxicant
gastrointestinal or liver toxicant
kidney toxicant
neurotoxicant
respiratory toxicant
skin or sense organ toxicant
More hazardous than most chemicals in 3 out of 10 ranking systems
On at least 8 federal regulatory lists

Phenoxyethanol: aka Antifreeze
EDF Suspected - developmental toxicant
reproductive toxicant
Less hazardous than most chemicals in 3 ranking systems

Sorbitol: EDF Suspected - gastrointestinal or liver toxicant
Less hazardous than most chemicals in 1 ranking system

Formaldehyde: EDF Recognized - carcinogen
Suspected - gastrointestinal or liver toxicant
immunotoxicant
neurotoxicant
reproductive toxicant
respiratory toxicant
skin or sense organ toxicant

More hazardous than most chemicals in 5 out of 12 ranking systems
On at least 8 federal regulatory lists
Ranked as one of the most hazardous compounds (worst 10%) to ecosystems and human health

Ask your pediatrician to name 3-4 ingredients in vaccines. If your pediatrician states "The amounts are so small" Ask them "How much bee venom does it take to cause anaphylaxis?"

Chemical Profiles and Definitions, visit www.scorecard.org to investigate chemical profiles Sources: EDF (Environmental Defense Fund) & MME (Mosby's Medical Encyclopedia)

"In some cases the cell lines that are used might be tumorigenic, that is, **they form tumors when injected into rodents.** Some of these tumor-forming cell lines may contain cancer-causing viruses that are not actively reproducing. Such viruses are hard to detect using standard methods. These latent, or "quiet," viruses pose a potential threat, since they might become active under vaccine manufacturing conditions."
FDA
<http://www.fda.gov/biologicsbloodvaccines/scienceresearch/biologicsresearchareas/ucm127327.htm>

ABORTED HUMAN FETAL TISSUE AND DNA

"Today, more than 23 vaccines are contaminated by the use of aborted fetal cells. There is no law that requires that consumers be informed that some vaccines are made using aborted fetal cells and contain residual aborted fetal DNA. While newer vaccines produced using aborted fetal cells do inform consumers, in their package inserts, that the vaccines contain contaminating DNA from the cell used to produce the vaccine, they do not identify the cells as being derived from electively aborted human fetuses."

The same group of physicians claims that there could be relations between this type of vaccines and diseases like diabetes, lupus, MS and autism:

"How could the contaminating aborted fetal DNA create problems? It creates the potential for autoimmune responses and/or inappropriate insertion into our own genomes through a process called recombination. There are groups researching the potential link between this DNA and autoimmune diseases such as juvenile (type I) diabetes, multiple sclerosis and lupus. Our organization, [3] Sound Choice Pharmaceutical Institute (SCPI), is focused on studying the quantity, characteristics and genomic recombination of the aborted fetal DNA found in many of our vaccines."

WOULD YOU RISK THIS WITH YOUR CHILD?

<http://www.physiciansforlife.org/content/view/1758/2/>