

# AGRONOMIC ALERT



## Corn Replanting Decisions in Northern and Central Illinois

Cool temperatures and wet conditions can put stress on corn germination and emerging seedlings. This may cause some concern about reduced corn stands. Before deciding to replant, evaluate the stand for population and uniformity. Next, compare yield potential of the existing stand with yield potential of the replant, as well as insurance or government program restrictions. Finally, if the decision is made to replant, consider various management practices discussed in this ALERT to help optimize yield potential.

### Evaluating the Existing Stand

When evaluating corn stands, only count plants that have a good chance of survival. Observe the uniformity of the stand across the field to decide whether replanting the entire field or portions of the field is justified. Spotting in without destroying the existing stand is not recommended<sup>3</sup>. There may be low areas in fields that will not survive, while the rest of the field will have a good stand. Replanting only the drowned out areas may be a viable option. Carefully evaluate the field for replant options.

One way to evaluate corn stands is to count the number of plants in a length of row equal to 1/1,000<sup>th</sup> of an acre based on row width (Table 1). Multiply the number of plants by 1,000 to get the plants per acre. Repeat the process in several locations in the field.

A more accurate method is to count 150 plants and measure the distance from start to finish with a measuring wheel. Divide the number of feet traveled into the appropriate factor in Table 2 to determine plant population. For example, if you walked 94 feet while counting 150 plants in 30-inch rows, the population is  $2,613,600 \div 94 = 27,804$ . Because a longer row length is counted, the samples are more representative and fewer locations are required.

Table 3. Percent of Maximum Yield Expected for Final Plant Stands Planted on Different Dates<sup>2</sup>.

Planting Date	Final Plant Population (x1000) Per Acre						
	10	15	20	25	30	35	40
	———— % of Maximum Yield ————						
April 1	54	68	78	88	95	99	99
April 10	57	70	81	91	97	100	100
April 20	58	71	81	91	97	100	99
April 30	58	70	80	89	95	97	96
May 9	55	68	77	86	91	93	91
May 19	50	63	72	80	85	86	84
May 29	44	56	65	73	77	78	75
June 8	35	47	56	63	67	67	64

Table 1. Stand count evaluation for 1/1,000<sup>th</sup> acre based on row width and number of plants in a given row length<sup>1</sup>.

Row Width (inches)	Row Length 1/1,000 <sup>th</sup> acre (feet, inches)
20	26' 2"
30	17' 5"
36	14' 6"
38	13' 9"

Table 2. Stand count evaluation factors, by row width, for measuring the distance when counting 150 plants<sup>1</sup>.

Row Width (inches)	Factor
20	3,920,400
30	2,613,600
36	2,178,000
38	2,063,350

Utilize the same factors for both 30-inch twin- and single-row stand counts; however, the plants in both the twin rows should be counted.

### Deciding Whether to Replant

After taking stand counts, consider the yield potential of the current stand, compared to the yield potential of the target replanting date and population, and the costs associated with replanting. Table 3 illustrates percent of maximum grain yield expected from various planting dates and final plant populations, based on uniform stands.

### When the Decision is Made to Replant

First you must remove the existing stand. Tillage is an option in many cases, and should be done at an adequate depth to properly control the original stand. Depending on the growth stage of the original stand, two tillage passes may be necessary. If a pre-emergence herbicide has been applied, tillage may decrease the efficacy of the herbicide by placing it deeper than ideal for satisfactory weed control. Additionally, tillage prior to replanting may result in loss of soil moisture. In drier areas, and where soil conditions allow, a more favorable option might be applying a herbicide to remove the original stand and then replanting without tillage. Roundup® brand agricultural herbicides are also an option when the original stand does not contain the Roundup Ready® Corn 2 trait or Roundup Ready® 2 Technology. If the original stand does contain the Roundup Ready® Corn 2 trait or Roundup

Continued on next page ▶

## Corn Replanting Decisions in Northern and Central Illinois

► Continued from page 1

Ready® 2 Technology, the tank mix options in Figure 1 can provide control of the original stand. The original plants must exhibit growing green tissue and 2 to 4 leaves for the herbicides to be effective. Complete kill may take up to 21 days after application.

Second, determine which relative maturity (RM) to use when replanting. As planting occurs after May 1, corn requires approximately 1.6 fewer growing degree days (GDDs) per day of delayed planting to reach flowering<sup>4</sup>. GDDs required to reach physiological maturity, or black layer, decreases approximately 6.8 GDDs per day of delayed planting after May 1<sup>4</sup>.

Third, decide on a management practice to protect against corn rootworm and other soil insects, as well as European Corn Borer (ECB). Most soil insecticides can not legally be applied twice in the same growing season in the same location in a field. Growers can replant over the old row and expect some control from the first application of soil insecticide. Another option is to use a different soil insecticide when replanting. A third option is to replant with corn containing the Genuity® family of traits. Depending on region and insect pests present, Genuity® SmartStax® RIB Complete™ and Genuity® VT Double PRO® RIB Complete™ corn blends, and Genuity® VT Triple PRO® Corn offer multiple modes of action for protection against a broad spectrum of insects. Later planted corn is more susceptible to second generation ECB. Research from The Ohio State University suggested significant advantages in overall yield and consistency of yield for corn with the corn borer-protection trait compared to their conventional counterparts, when planted in late May and early June<sup>3</sup>.

Finally, another consideration is later planted corn also has a greater chance of being exposed to heat and drought stress during pollination. This risk can be managed by selecting corn with heat and drought tolerance and early flowering.

Sources: <sup>1</sup> Nafziger, E.D. et al. 2002. *Illinois agronomy handbook*, 23rd edition. U of I Printing Services. Urbana, IL. 27-33.

### Figure 1. Tank Mix Options for Removing Corn for Replant

#### Option 1:

Select Max®	6 oz/acre
Non-ionic Surfactant	0.25% v/v
Ammonium Sulfate	2.5 to 4 lbs./acre

- Apply in a minimum of 10 gallons of water per acre.
- Treat prior to volunteer corn reaching 12 inches in height.
- Replant corn no sooner than 6 days after application.
- May be applied as a tank mixture with Roundup® brand agricultural herbicides.
- Care must be taken to avoid in-field boom (spray) overlaps or excessive crop injury may occur.

#### Option 2:

Gramoxone Inteon®	2.5 pt/acre (for corn 1 to 3" tall) OR 3 pt/acre (for corn 3 to 6" tall)
	plus
Metribuzin DF®	3 oz/acre
	plus
COC	1% v/v

- Apply in a minimum of 20 gallons of water per acre.
- There are no plant back restrictions.
- Clarity® herbicide may be added at 8 oz/acre for enhanced control of marestail and other tough-to-control broadleaves.
- 2,4-D at 1 pt/acre may be added, but typically requires a 7 day plant back interval.
- Check the Clarity and 2,4-D labels for specific instructions.

<sup>2</sup> Nafziger, E.D. et al. 2009. *Illinois agronomy handbook*, 24th edition. University of Illinois Board of Trustees. Urbana, IL. 24.

<sup>3</sup> Thomison, P. 2005. *Replant considerations: hybrid selection issues*. The Ohio State Univ. C.O.R.N. Newsletter 2005-13. <http://corn.osu.edu> (verified 4/11/2012)

<sup>4</sup> S. Brouder et al. 2007. *Corn & Soybean Field Guide*. Purdue University

Monsanto Company is a member of Excellence Through Stewardship® (ETS). Monsanto products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Monsanto's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for this product. Excellence Through Stewardship® is a registered trademark of Biotechnology Industry Organization. **BT products** may not yet be registered in all states. Check with your Monsanto representative for the registration status in your state. **IMPORTANT IRM INFORMATION: RIB Complete™** corn does not require the planting of a structured refuge **except** in the Cotton-Growing Area where corn earworm is a significant pest. **Genuity® SmartStax® RIB Complete™ and Genuity® VT Double PRO® RIB Complete™** corn are blended seed corn products. See the IRM/Grower Guide for additional information. Always read and follow IRM requirements. **Individual results may vary**, and performance may vary from location to location and from year to year. This result may not be an indicator of results you may obtain as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Roundup Ready® crops contain genes that confer tolerance to glyphosate, the active ingredient in Roundup® brand agricultural herbicides. Roundup® brand agricultural herbicides will kill crops that are not tolerant to glyphosate. **Tank mixtures:** The applicable labeling for each product must be in the possession of the user at the time of application. Follow applicable use instructions, including application rates, precautions and restrictions of each product used in the tank mixture. Monsanto has not tested all tank mix product formulations for compatibility or performance other than specifically listed by brand name. Always predetermine the compatibility of tank mixtures by mixing small proportional quantities in advance. Genuity and Design®, Genuity Icons, Genuity®, RIB Complete™, Roundup Ready 2 Technology and Design®, Roundup Ready®, Roundup®, SmartStax®, VT Double PRO®, and VT Triple PRO® are trademarks of Monsanto Technology LLC. Leaf Design™ is a servicemark of Monsanto Company. Ignite® and LibertyLink® and the Water Droplet Design® are registered trademarks of Bayer. Herculex® is a registered trademark of Dow AgroSciences LLC. Select Max® is a registered trademark of Valent U.S.A. Corporation. Respect the Refuge and Corn Design® and Respect the Refuge® are registered trademarks of National Corn Growers Association. All other trademarks are the property of their respective owners. ©2012 Monsanto Company. 04162012EJP



Before opening a bag of seed, be sure to read, understand and accept the stewardship requirements, including applicable refuge requirements for insect resistance management, for the biotechnology traits expressed in the seed as set forth in the Monsanto Technology/Stewardship Agreement that you sign. By opening and using a bag of seed, you are reaffirming your obligation to comply with the most recent stewardship requirements.

