

Mycobacterial Disease Other Than Tuberculosis (MOTT)

REPORTING INFORMATION

- **Class B:** Report the case, suspected case, and/or a positive laboratory result to the county local health department where the patient resides by the close of the next business day. If patient residence is unknown, report to the local public health department in which the reporting health care provider or laboratory is located.
- Health care providers and laboratories report using Form(s) and/or Mechanism: [Ohio Confidential Reportable Disease form](#) (HEA 3334, rev. 1/09), [Positive Laboratory Findings for Reportable Disease form](#) (HEA 3333, rev. 8/05), the Ohio Disease Reporting System (ODRS), Electronic Disease Reporting (ELR), or telephone.
- Local public health departments report the case, suspected case and/or a positive laboratory result to the Ohio Department of Health (ODH) via the Ohio Disease Reporting System (ODRS) by the end of the next business day.
- Key fields for ODRS reporting include: laboratory date collected, specimen type, test name, result, organism, and result date.

AGENTS

There are hundreds of mycobacteria that may be pathogenic to humans. MOTT includes all mycobacteria other than those belonging to *M. tuberculosis* complex (i.e. *M. tuberculosis*, *M. bovis*, *M. bovis BCG*, *M. africanum*, *M. microti*, *M. canetti*, *M. pinipedi*) and *M. leprae*, which causes Hansen Disease ([leprosy](#)).

CASE DEFINITION (ODH, 2009)

The Centers for Disease Control and Prevention has not established a case definition for MOTT. Reports should be based upon the clinical signs and symptoms and the laboratory criteria described below.

Clinical Description

A bacterial infection caused by mycobacteria other than those belonging to *M. tuberculosis* complex and *M. leprae*, other separately reportable Class B diseases.

Laboratory Criteria for Diagnosis

- Isolation of a MOTT from a clinical specimen *or*
- Demonstration of a MOTT from a clinical specimen by nucleic acid amplification

Case Classification

Suspect: A clinically compatible case without laboratory confirmation.

Confirmed: A clinically compatible case that is laboratory confirmed.

Not a Case: This status will not generally be used when reporting a case, but may be used to reclassify a report if investigation revealed that it was not a case.

SIGNS AND SYMPTOMS

Indications of a MOTT infection will vary depending on the MOTT and the part of the body infected.

DIAGNOSIS

The definitive diagnosis of MOTT requires the isolation of a MOTT from the patient. MOTT may infect any part of the body.

EPIDEMIOLOGY

Source

MOTT are found in humans, animals, and the environment.

Occurrence

Present in all parts of the world.

Mode of Transmission

Dependent upon the MOTT involved; transmission may be person-to-person, animal-to-person, or environmental.

Period of Communicability

Dependent upon the MOTT involved.

Incubation Period

Dependent upon the MOTT involved.

PUBLIC HEALTH MANAGEMENT

Case

Management of MOTT infection is handled in the private sector; public health is not involved unless the person is also infected with *M. tuberculosis* complex or *M. leprae*.

Prevention and Control

Dependent upon the MOTT involved; may involve respiratory and contact precautions.