

Psychological Medicine

<http://journals.cambridge.org/PSM>


Additional services for *Psychological Medicine*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Letter to the Editor: Childhood trauma may combine synergistically with stimulant use rather than cannabis use to predict psychosis

MICHAEL DALY

Psychological Medicine / Volume 42 / Issue 02 / February 2012, pp 445 - 446
DOI: 10.1017/S0033291711002546, Published online: 16 November 2011

Link to this article: http://journals.cambridge.org/abstract_S0033291711002546

How to cite this article:

MICHAEL DALY (2012). Letter to the Editor: Childhood trauma may combine synergistically with stimulant use rather than cannabis use to predict psychosis. *Psychological Medicine*, 42, pp 445-446 doi:10.1017/S0033291711002546

Request Permissions : [Click here](#)

Correspondence

Psychological Medicine, 42 (2012).

doi:10.1017/S0033291711002546

First published online 16 November 2011

Letter to the Editor

Childhood trauma may combine synergistically with stimulant use rather than cannabis use to predict psychosis

Early experiences of physical and sexual abuse are considered to contribute to the development of psychosis later in life (Read *et al.* 2005; Schreier *et al.* 2009; Arseneault *et al.* 2011). Similarly, cannabis use has been linked to an increased risk of psychosis and an earlier age of onset of psychotic illness (Moore *et al.* 2007; Large *et al.* 2011). A growing number of studies have shown that childhood trauma and cannabis use can combine more-than-additively to produce a highly elevated risk of psychotic symptoms and psychosis (e.g. Houston *et al.* 2008; Harley *et al.* 2010).

Most recently, two studies published in *Psychological Medicine* have shown that: (i) psychotic symptoms are particularly likely to occur amongst cannabis users with a history of childhood physical or sexual mistreatment (Konings *et al.* 2011), and (ii) that psychosis risk is enhanced amongst cannabis users who have experienced non-consensual sex before the age of 16 years (Houston *et al.* 2011). Crucially, neither study considered the possibility that the risk posed by prior trauma could be enhanced amongst cannabis users because individuals in this group are also likely to have taken illicit psychostimulants implicated in the development of psychosis (Fergusson & Horwood, 2000; Lynskey *et al.* 2003; Curran *et al.* 2004; Barnett *et al.* 2007). To test this idea, I examined data from 7125 participants drawn from the Adult Psychiatric Morbidity Survey 2007, as utilized in Houston *et al.* (2011).

As shown in Houston *et al.* (2011), cannabis use moderated the link between non-consensual sex in childhood and psychotic disorder in the last week [odds ratio (OR) 10.53, 95% confidence interval (CI) 1.14–99.64; all analyses were adjusted for age, gender, ethnicity, education, employment, alcohol use, sexual trauma after age 16 years, presence of neurotic disorder, and, where appropriate, cannabis and stimulant use]. Similarly, stimulant use (cocaine, ecstasy or amphetamines) combined with non-consensual sex before the age of 16 years to predict a raised risk of psychosis (OR 16.75, 95% CI 1.79–157.2). Cannabis dependency levels did not interact with

non-consensual sex to predict psychosis, suggesting that psychostimulant use is not merely a proxy for the effects of high levels of cannabis consumption.

Critically, adjusting for the under-16 sex × stimulant use interaction removed the link between the under-16 sex × cannabis use interaction and psychosis (OR 8.22, 95% CI 0.48–140.9). As anticipated, the link between under-16 sex and psychosis amongst cannabis users (OR 17.43, 95% CI 2.61–116.43; illustrated in Fig. 1a) was found to be attributable to a strong link between non-consensual sex in childhood and psychosis amongst the 37.1% of cannabis users who have also taken stimulants (OR 77.7, 95% CI 10.39–581.1 in age and gender adjusted analysis; OR 70.71, 95% CI 1.38–3631 in fully adjusted analysis), as shown in Fig. 1b. Early non-consensual sex was unrelated to psychosis amongst those who had used cannabis alone.

Taken together, these findings suggest that stimulant use amongst cannabis users rather than cannabis use alone may enhance the impact of childhood trauma on the likelihood of developing psychotic disorder. Future studies examining cannabis × trauma interactions in psychosis should include the main effect of stimulants and interactions between trauma and stimulants in their analyses. Such an approach may uncover novel relations and will ensure that those examining synergistic relations between cannabis use and adverse experiences accurately identify the specific drugs that contribute to psychosis risk.

Declaration of Interest

None.

References

- Arseneault L, Cannon M, Fisher HL, Polanczyk G, Moffitt TE, Caspi A (2011). Childhood trauma and children's emerging psychotic symptoms: a genetically sensitive longitudinal cohort study. *American Journal of Psychiatry* 168, 65–72.
- Barnett JH, Werners U, Secher SM, Hill KE, Brazil R, Masson K, Pernet DE, Kirkbride JB, Murray GK, Bullmore ET, Jones PB (2007). Substance use in a population-based clinic sample of people with first-episode psychosis. *British Journal of Psychiatry* 190, 515–520.
- Curran C, Byrappa N, McBride A (2004). Stimulant psychosis: systematic review. *British Journal of Psychiatry* 185, 196–204.




Fig. 1. Relationship between non-consensual sex before the age of 16 years and psychosis for: (a) those who have used cannabis and those who have not taken cannabis or stimulants (cocaine, amphetamines or ecstasy); and (b) cannabis users who have taken stimulants, cannabis-only users and those who have not taken cannabis or stimulants. Note: participants who have used stimulants but not cannabis (approximately 1% of the sample) were excluded for illustration purposes.

Fergusson DM, Horwood LJ (2000). Does cannabis use encourage other forms of illicit drug use? *Addiction* **95**, 505–520.

Harley M, Kelleher I, Clarke M, Lynch F, Arseneault L, Connor D, Fitzpatrick C, Cannon M (2010). Cannabis use and childhood trauma interact additively to increase the risk of psychotic symptoms in adolescence. *Psychological Medicine* **40**, 1627–1634.

Houston JE, Murphy J, Adamson G, Stringer M, Shevlin M (2008). Childhood sexual abuse, early cannabis use and psychosis: testing an interaction model based on the National Comorbidity Survey. *Schizophrenia Bulletin* **34**, 580–585.

Houston JE, Murphy J, Shevlin M, Adamson G (2011). Cannabis use and psychosis: revisiting the role of childhood trauma. *Psychological Medicine* **41**, 2339–2348.

Konings M, Stefanis N, Kuepper R, de Graaf R, ten Have M, van Os J, Bakoula C, Henquet C (2011). Replication in two independent population-based samples that childhood maltreatment and cannabis use synergistically impact on psychosis risk. *Psychological Medicine*. Published online: 15 June 2011. doi:10.1017/S0033291711000973.

Large M, Sharma S, Compton MT, Slade T, Nielsen O (2011). Cannabis use and earlier onset of psychosis: a systematic meta-analysis. *Archives of General Psychiatry* **68**, 555–561.

Lynskey MT, Heath AC, Bucholz KK, Slutske WS, Madden PAF, Nelson EC, Statham DJ, Martin NG (2003). Escalation of drug use in early-onset cannabis users vs

co-twin controls. *Journal of the American Medical Association* **299**, 427–433.

Moore TH, Zammit S, Lingford-Hughes A, Barnes RE, Jones PB, Burke M, Lewis G (2007). Cannabis use and risk of psychotic or affective mental health outcomes: a systematic review. *Lancet* **370**, 319–328.

Read J, van Os J, Morrison AP, Ross CA (2005). Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica* **112**, 330–350.

Schreier A, Wolke D, Thomas K, Horwood J, Hollis C, Gunnell D, Lewis G, Thompson A, Zammit S, Duffy L, Salvi G, Harrison G (2009). Prospective study of peer victimization in childhood and psychotic symptoms in a nonclinical population at age 12 years. *Archives of General Psychiatry* **66**, 527–536.

MICHAEL DALY^{1,2,3}

¹ Health Services Research Unit, Health Sciences Building, University of Aberdeen, Foresterhill, Aberdeen UK

² UCD Geary Institute, University College Dublin, Belfield, Dublin 4, Republic of Ireland

³ School of Psychological Sciences, University of Manchester, Oxford Road, Manchester, UK

Address for correspondence:

Dr Michael Daly
Health Services Research Unit,
Health Sciences Building, University of Aberdeen,
Foresterhill, Aberdeen AB25 2ZD, UK.
(Email: m.daly@abdn.ac.uk)